



**Jonathan Green**

# VERI-GREEN®

## Lawn Food with Crabgrass Preventer

**PREVENTS CRABGRASS & LISTED WEEDS & GRASSES** | **FEEDS YOUR LAWN FOR UP TO 3 MONTHS** | **SEASON LONG CRABGRASS CONTROL**

**CONTROL CRABGRASS BEFORE IT EMERGES... AND UP TO 4 WEEKS AFTER IT EMERGES!**

**20-0-3**

**ACTIVE INGREDIENT:**  
 Dithiopyr, 3,5-pyridinedicarboxylic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-5,5-dimethyl ester ..... 0.125%  
 99.875%  
**OTHER INGREDIENTS:** ..... 100.000%  
 TOTAL: ..... 100.000%

EPA REG. NO. 62719-483-50931  
 EPA EST. NO. 075527-PA-12  
 90561-NC-001 89135-AZ-001  
 09243-CF-1 8378-IN-1  
 69759-OR-2 69759-OR-004 (LOT BEGINS BAW)  
 Code on bag indicates manufacturing location.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
 Refer to back of bag for Directions for Use

• Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.  
 • Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.  
 • For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, do not exceed 4.6 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre). In the state of New York, this product may be applied at a minimum of 2.4 lb per 1000 sq ft per year (equivalent to 0.128 lb active ingredient per acre).

GUARANTEED ANALYSIS	
Total Nitrogen (N) .....	20%
20.00% Water Soluble Nitrogen*	
Soluble Potash (K <sub>2</sub> O) .....	3%
Derived From: Polymer Coated Urea, Urea, Muriate of Potash.	
*5.0% controlled release urea nitrogen from polymer coated urea	
Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.aapfco.org/metals.html">http://www.aapfco.org/metals.html</a>	

**FIRST AID**  
**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, the give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**HOT LINE NUMBER**  
 Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**  
 Causes Eye Irritation • Harmful if Inhaled  
 Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

**ENVIRONMENTAL HAZARDS**  
 This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.  
**Notice:** Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.  
 In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
**STORAGE AND DISPOSAL**  
 Do not contaminate water, foodstuffs, feed or seed by storage or disposal.  
**Pesticide Storage:** Store this product only in its original container in a dry, cool, secured storage area.  
**Container Handling:** Nonrefillable container. Do not reuse or refill this container.  
**If empty:** Place in trash or after recycling if available.  
**If partially filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**USE DIRECTIONS FOR TURF**  
 This turf fertilizer product, containing Dimension® herbicide, can be applied 4 weeks after preemergent crabgrass timing and provides up to 4 months of crabgrass control.  
 This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only, but is effective up to 4 weeks after crabgrass has germinated. This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after crabgrass germination.  
**Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:**

COOL-SEASON GRASSES:			WARM-SEASON GRASSES:		
Bentgrass, creeping	Fescue, fine	Ryegrass, perennial	Bahiagrass	Buffalograss	Centipedeagrass
Bluegrass, Kentucky	Fescue, tall		Bermudagrass	Carpetgrass	St. Augustinegrass
					Zoysiagrass

**USE PRECAUTIONS FOR TURF**  
 This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. **Exception:** In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.  
 • Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.  
 • For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aeration, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.  
 • **Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 1-6 weeks after a split application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

**USE RESTRICTIONS**  
 • Do not use clippings from treated turf for mulching around vegetables or fruit trees.  
 • Do not allow people or pets to enter treated area until dust has settled.

**APPLICATION TIMING**  
**SPRING APPLICATION**  
**Control of Crabgrass:** This product provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when crabgrass plants begin to tiller (spread). Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.  
**Control or Suppression of Other Grass and Broadleaf Weeds:** For control or suppression of other grass and broadleaf weeds, this product must be applied as a preventative treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

**FALL APPLICATION**  
 This product can also be applied from late summer through early fall (late August through November). When in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig turfgrasses within 12 weeks after application.

**TIPS FOR IMPROVED WEED CONTROL**  
 • For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.  
 • The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

### WEEDS CONTROLLED OR SUPPRESSED

This product will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior their germination.

GRASSES		BROADLEAF WEEDS	
barley	goosegrass	bittercrack †	medic, black
barnyardgrass	foxtail, green	carpetweed	mustard
bluegrass, annual	foxtail, yellow	chickweed †	oxalis, buttercup
"lame"	kikuygrass †	geranium, Carolina †	pineappleweed
crabgrass, large	oats, wild	henbit	pigweed, redroot
crabgrass, smooth	ryegrass, (annual & perennial)	knotted, prostrate	parsleywort †
crabgrass, Southern	"perennial"	lespedeza, common †	purslane, common
crowfootgrass †	sandbar	marestail	rocket, London
dallisgrass	smulgrass		

† Suppression only

### LENGTH OF CONTROL

This product will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Refer to the map below to determine the region of the U.S. that includes your location. Use the table below to determine the amount of this product to apply in your location based on the desired length of weed control or need for postemergence crabgrass control.

Use "Regions" within the United States:

Region of U.S.	Amount of this Product to Apply (Pounds of product to apply per 1,000 square feet of turfgrass)		
	2-3 Months	3-4 Months	Early Postemergence Control
North	2.4 lb/1000 sq ft (0.128 lb ai/acre)	3.2 lb/1000 sq ft (0.174 lb ai/acre)	4.0 - 9.2 lb/1000 sq ft † (0.22 - 0.5 lb ai/acre)
South	3.5 lb/1000 sq ft (0.192 lb ai/acre)	4.0 - 4.7 lb/1000 sq ft † (0.22 - 0.256 lb ai/acre)	4.0 - 9.2 lb/1000 sq ft † (0.22 - 0.5 lb ai/acre)

1 1,000 square feet equals an area 100 feet X 10 feet or square area of about 32 feet on a side.  
 † Use the higher rate in rate range in areas of high weed pressure or for increased length of control.  
**Maximum Application Rates:**  
 • Do not apply more than 6 times per year and do not apply more than 9.2 lb of this product per 1000 sq ft (0.5 lb ai/acre) per application.  
 • Do not apply more than 27.6 lb of this product/1000 sq ft per year (equivalent to 1.5 lb of active ingredient per acre, respectively).  
 • In the state of New York, do not exceed 4.6 lb per 1000 sq ft per year of this product (equivalent to 0.25 lb of active ingredient per acre).  
 • To minimize the potential for plant injury, do not make additional applications within five weeks of previous application.

### APPLICATION INSTRUCTIONS AND EQUIPMENT

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the specified application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area.  
 More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.  
 Check operation frequently to verify calibration and proper functioning.  
 \*These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product.

### CALIBRATION SETTINGS AT SPECIFIED RATE

BROADCAST (ROTARY) SPREADERS	3.2 lb, 1,000 sq. ft.	DROP SPREADERS	3.2 lb, 1,000 sq. ft.
Jonathan Green Standard	5-1/2	Central	6
Deluxe & Pro Hand Held	14	EZ Republic (Ortho)	4
AgriFab	4	Earthway (Ames)	14
Chapin 81000A	5	Precision (Red Devil)	5
Cyclone	4-1/2	Scotts Drop**	7
EZ Republic (Ortho)	3		
Earthway (Ames)	14		
Lesco	11 - 12/D - E		
Precision (Red Devil)	5		
Scotts Rotary*	5		
Scotts EasyGreen	26		

\*\*Includes Accugreen 1000 - 2000 & Classic  
 \*Includes EdgeGuard, Elite, Standard, Basic, LawnPro & Speedy Green 1000 - 2000

**TERMS AND CONDITIONS OF USE**  
 If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

**WARRANTY DISCLAIMER**  
 Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**INHERENT RISKS OF USE**  
 It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

**LIMITATION OF REMEDIES**  
 To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:  
 1. Refund of purchase price paid by buyer or user for product bought, or  
 2. Replacement of amount of product used.  
 To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.  
 The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

**Dimension** Specialty Herbicide  
 Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**MADE IN USA**

